



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

If  $n$  be an odd No.  $\beta_{\frac{1}{2}(n+1)}$  will be the middle term and its mean value will be  $\frac{1}{2}\alpha$ . If  $n$  be an even No.  $\beta_{\frac{1}{2}n}$  and  $\beta_{\frac{1}{2}(n+1)}$  are middle terms; their mean values are  $\frac{1}{2}\alpha + \frac{\frac{1}{2}\alpha}{n+1}$  and  $\frac{1}{2}\alpha - \frac{\frac{1}{2}\alpha}{n+1}$ , and the mean of these values is  $\frac{1}{2}\alpha$ . The sum of  $n$  values is  $\frac{1}{2}an$ , and hence the mean of all the values is  $\frac{1}{2}\alpha$ .

Suppose now we take the values of the three tickets, as estimated per question by the independent voters, and that these values are as the Nos. 1, 2, 3. Then, substituting  $n$  by 3,  $\alpha$  by 100, and  $r$  severally by 1, 2, 3, we shall have for the mean value of the most worthy ticket,  $\frac{3}{4} \times 100 = 75$ ,  
 for the mean value of the next best,  $\frac{2}{4} \times 100 = 50$ ,  
 for the mean value of the third “ ,  $\frac{1}{4} \times 100 = 25$ ;  
 while the *mean* of all the values will be  $\frac{1}{3} \times \frac{3}{2} \times 100 = 50$ .

---

### SOLUTION OF A PROBLEM.

---

BY WILLIAM HOOVER, SUPERINTENDENT OF SCHOOLS, WAPAKONETA, O.

*Problem.*—A triangle is formed by joining the centers of the escribed circles of a triangle; the same is done for the new triangle, and so on *ad infinitum*. Show that the final triangle is equilateral.

*Solution.*—If  $A$  be one of the angles of the given triangle, and  $A_1, A_2, A_3$ , &c., the successive corresponding angles, we have  
 $A_1 = (\frac{1}{2})(\pi - A)$ ,  $A_2 = (\frac{1}{2})^2(\pi + A)$ ,  $A_3 = (\frac{1}{2})^3(3\pi - A)$ ,  $A_4 = (\frac{1}{2})^4(5\pi + A)$ .

The fraction for the  $n$ th angle is  $(\frac{1}{2})^n$ , the coefficient of  $\pi$  is the general term of the recurring series 1, 1, 3, 5, &c.,  $\frac{1}{3}[2^n + (-1)^{n-1}]$ , and of  $A$ , it is  $(-1)^n$ . Hence

$$A_n = \frac{1}{2^n} \left[ \frac{1}{3}\pi[2^n + (-1)^{n-1}] + (-1)^n A \right].$$

When  $n$  is infinite  $A_n = \frac{1}{3}\pi$ , and the triangle is equilateral.

---

### SOLUTION OF PROB. 282 WITH CONDITION (3) CHANGED.

---

BY CHAS. H. KUMMELL, U. S. LAKE SURVEY.

The solution of this problem (see pp. 25, 26) is in accordance with the given conditions. However it cannot be admitted, as assumed in condition (3), that it is equally probable that only one loom should be out of order as

that any other number should be out of order. I propose to solve the problem by assuming  $a$  looms, each netting  $m$  dollars a day when in order,  $x$  weavers, each getting  $n$  dollars a day, whether working or not, and  $p$  the probability of a single loom being out of order. We have then to take the general mean of the following cases:

No. of looms   Profits.   Probability.  
in order.

|       |                 |   |             |
|-------|-----------------|---|-------------|
| $a$   | $mx$            | $(1-p)^a$   | $= P_a$     |
| $a-1$ | $mx$            | $\frac{a}{1}(1-p)^{a-1} p$  | $= P_{a-1}$ |
| $a-2$ | $mx$            | $\frac{a}{1} \cdot \frac{a-1}{2} (1-p)^{a-2} p^2$                               | $= P_{a-2}$ |
| .     | .               | .   | .           |
| $x$   | $mx$            | $\frac{a}{1} \cdot \frac{a-1}{2} \dots \frac{x+1}{a-x} (1-p)^x p^{a-x}$         | $= P_x$     |
| $x-1$ | $mx-(m+n)$      | $\frac{a}{1} \cdot \frac{a-1}{2} \dots \frac{x}{a-x+1} (1-p)^{x-1} p^{a-x+1}$   | $= P_{x-1}$ |
| $x-2$ | $mx-2(m+n)$     | $\frac{a}{1} \cdot \frac{a-1}{2} \dots \frac{x-1}{a-x+2} (1-p)^{x-2} p^{a-x+2}$ | $= P_{x-2}$ |
| .     | .               | .   | .           |
| 1     | $mx-(x-1)(m+n)$ | $\frac{a}{1} (1-p) p^{a-1}$   | $= P_1$     |
| 0     | $mx-x(m+n)$     | $p^a$   | $= P_0$     |

We have then the average profit

$$y_x = mx - (m+n) \sum_1^x [sP_{x-s}].$$

If this is a maximum then

$y_x - y_{x-1} = \Delta y_x = m - (m+n) \sum_1^x [P_{x-s}]$  is small and positive,  
&  $y_{x+1} - y_x = \Delta y_{x+1} = m - (m+n) \sum_1^{x+1} [P_{x+1-s}]$  is small and negative,

Placing  $a = 60$ ,  $m = 3$ ,  $n = 1$  and  $p = 0.1$  we find  $\Delta y_{56} = + 0.021$  and  $\Delta y_{57} = - 0.113$ . Therefore  $x = 56$  weavers.

## ANSWERS TO QUERIES IN NUMBER FOUR.

“QUERY BY PROF. A. HALL.—(1). If  $v$  be the potential function we have the equation given by Laplace

$$\frac{d^2 v}{dx^2} + \frac{d^2 v}{dy^2} + \frac{d^2 v}{dz^2} = 0,$$

which holds for a point outside the attracting body. For a point inside this body we have the equation given by Poisson,